

U. S. DEPARTMENT OF COMMERCE  
ENVIRONMENTAL SCIENCE SERVICES ADMINISTRATION  
WEATHER BUREAU

WEATHER BUREAU TECHNICAL MEMORANDUM SR-37

THE RELATIONSHIP OF K-VALUES TO PROBABILITY OF SHOWERS IN THE MID-SOUTH  
Kenneth E. Bryan  
WBAS, Memphis, Tennessee

SOUTHERN REGION HEADQUARTERS  
SCIENTIFIC SERVICES DIVISION  
TECHNICAL MEMORANDUM NO. 37

FORT WORTH, TEXAS  
October 1967





# THE RELATIONSHIP OF K-VALUES TO PROBABILITY OF SHOWERS IN THE MID-SOUTH

Kenneth E. Bryan  
WBAS, Memphis, Tennessee

## ABSTRACT

This article describes and illustrates a forecasting technique used at Memphis for shower and thundershower potential for a given day during the summer months over the Mid-South agricultural area. This technique is related to the precipitation probability now currently used by the ESSA Weather Bureau. The statistical analysis used in this development is shown.

## INTRODUCTION

Accurate forecasting of showers and thundershowers from late spring until early fall is one of the most important services for the Mid-South agricultural program; however, achieving a high degree of accuracy is a very significant problem. Air mass thundershowers, as well as occasional frontal showers and thundershowers, occur during this time with air mass activity usually playing the dominant role.

In the absence of early machine prog analysis, K-values have proved to be a very quick and useful tool. We concede that the machine can organize data more quickly and consider more factors, but the K-values have considerable merit. The K-chart is a factor to analyze in conjunction with other information, but overall it gives one a better idea where to look and a little better visual analysis.

## PROCEDURE OF INVESTIGATION

The forecast problem was defined as: What percent of the Mid-South agricultural area including the Missouri Bootheel, West Tennessee, Northern Half of Mississippi, Louisiana Delta, and Arkansas will receive measurable rain during the 24-hour period following the 1200Z observational data? (See Fig. 1)

Rainfall records from 75 reporting stations connected by the agricultural and public service communications network over the Mid-South were used to determine a percent probability of these stations receiving measurable rain during this 24-hour period.

K-values, as described by J. J. George, combine both moisture and stability parameters into a single, easily computed index. (See Fig. 2) These values can be computed for the areas east of the Rocky Mountains in approximately 10 minutes. Wind data at 700 MB is also plotted for advection information. The chart is then analyzed by drawing K-value isopleths with intervals of five units. (Refer to Fig. 3) Rainfall for the following 24-hour period is shown in the lower right hand corner of the figure. This simple K-chart has proved very useful in the agricul-

tural program as well as the state and aviation forecast programs, and has led to this attempt to objectively relate K-values to the point probability of showers and thundershowers. A Chi-Square analysis shows statistical significance at the 99 percent confidence level between increments of rainfall amount and K-value amount.

Figure 4 shows a breakdown of total cases of measurable rainfall for 75 stations in the Mid-South for the summer months from May through September distinguishing them from precipitation amounts and K-values. Figure 5 portrays the graphical picture showing percent of total of each K-value for various increments of precipitation amounts. Here it is seen that there are more cases of measurable precipitation with increasing K-values.

By taking a portion of cases and graphically portraying them, it is found with precipitation values between .50 and 1.99 inches of rain that the percent of total cases more than doubles when equal or greater than 30K is used and that there is slightly over 90 percent of total cases when equal or greater than 20K is used. (See Figure 6A) When using one-quarter inch or more rain in comparison with .50 to 1.99 inches the percent of total cases increases slightly for equal or greater than 30K and also for equal or greater than 20K. (Refer to Figure 6B) A total of 10,857 cases of rain and no rain was used in this study, 4441 of no rain cases with less than 20K, 2602 cases of no rain with 20-29K, and 1390 cases of no rain with 30K or greater.

Figure 6C shows the probability of rain for any particular station over the Mid-South. Cases with rain over total cases (rain and no rain) give the percent probability rounded to the nearest increment of 10. As would be expected, the higher the K-value the greater the probability. It is also found that 4 percent probability of rain or 96 percent probability of no rain is noted for cases less than 20K. Of course 4 percent rounded to the nearest increment of 10 is zero.

#### SUMMARY AND CONCLUSIONS

A definite relationship of K-values to shower and thundershower occurrence has been established in the Mid-South. This addition of K-values as a tool in forecasting point probability of showers and thundershowers has definitely improved this type of forecast to the extent that all users contacted express satisfaction with this service. Not only does this study work well with air mass activity but also can be applied with good accuracy to approaching cold fronts.

The main advantage of this technique is that it can be prepared early around 7 a.m., some two hours before remaining raob data and computer progs are received.

ACKNOWLEDGMENT

The author wishes to thank Mr. Russ Alderman for his statistical help during the preparation of this paper, and the rest of the agricultural forecasters and others who have helped collect data since 1965.

BIBLIOGRAPHY

1. J. J. George, "Weather Forecasting for Aeronautics" 1960, pp. 409-415.
2. Jack Hollis and Kenneth E. Bryan, "The Relationship of K-Values to Areal Coverage of Showers in the Mid-South", Technical Memorandum No. 2, August 1965.



WBAS MEMPHIS AREA FOR WHICH  
AGRICULTURAL FORECASTS ARE ISSUED





FIGURE 1

K = (850 MB Temp. + Dewpoint) - (700 MB Spread) - (500 MB Temp.)

UXUS 30 301250					
FTW					
GSW GW TT 72259 12 10196 85569 20082 02706 70201 05523 01016					= 30 K
50589 58507 00903 020					
LIT TT 72340 12 10196 85558 19502 03506 70195 06014 00212 50588					= 20 K
57797 03013 090					
SHV TT 72248 12 10196 85553 17518 00113 70183 08662 00106 50587					
59839 03404 080					

FIGURE 2





K-CHART AND FOLLOWING 24 HR. RAINFALL  
FIGURE 3

# K VALUES

	≤ 14	15 - 19	20 - 24	25 - 29	30 - 34	35 & >
0.01 - .09	37	27	108	167	276	134
.10 - .24	15	18	47	110	211	102
.25 - .49	9	15	54	97	174	134
.50 - .99	18	26	43	87	162	108
1.00 - 1.99	5	12	33	46	85	69
≥ 2.00	0	5	3	13	20	25

## PRECIPITATION

## CASES

FIGURE 1.




FIGURE 5

.50 - 1.99 inches

1/4 inch or more




FIGURE 6A

FIGURE 6B

FIGURE 6C

LIST OF SOUTHERN REGION TECHNICAL MEMORANDA

- No. 1 Selection of Map Base for Small Scale Analysis and Prediction. Woodrow W. Dickey - May 19, 1965
- No. 2 The Relationship of K-Values to Areal Coverage of Showers in the Mid-South. Jack Hollis and Kenneth E. Bryan - August 1965
- No. 3 Some Notes on Waterspouts around the Lower Keys. R. Larry Mayne - August 1965
- No. 4 Tornadoes Associated with Cyclones of Tropical Origin - Practical Features. E. L. Hill, William Malkin and W. A. Schulz, Jr. - September 1965
- No. 5 Radar Echoes Associated with Waterspout Activity. Dorus D. Alderman - October 1965
- No. 6 Probability Forecasting. Woodrow W. Dickey - October 1965
- No. 7 Short Period Forecasting. Jeter A. Pruett - November 1965
- No. 8 Southwest Texas Soaring Weather. David H. Owens - November 1965
- No. 9 A Survey of Research in Agricultural Meteorology. Donald A. Downey - November 1965
- No. 10 A Quick Look at the Results of One Month's Precipitation Probability Forecasting. George T. Gregg - January 1966
- No. 11 Severe Storm Warning Systems in the Southern Region. Staff Members of Operations Division, Southern Region - February 1966
- No. 12 The Lubbock Snowstorm of February 20, 1961. Billie J. Cook - April 1966
- No. 13 Summary of Probability of Precipitation Forecasts in the Southern Region for the Period January through March 1966. Woodrow W. Dickey - May 1966
- No. 14 Air Pollution Meteorology and Transport of Pollutants. Jose A. Colon - June 1966
- No. 15 On the Mechanisms for the Production of Rainfall in Puerto Rico. Jose A. Colon - June 1966
- No. 16 Teletype Techniques and Presentation Procedures for Public Weather Circuits. Jack Riley - June 1966
- No. 17 Summary of the Pre-FP - Post-FP Forecast Verification Experiment. Woodrow W. Dickey - June 1966
- No. 18 Fire Weather in the Southeast. Robert A. Mitchem - July 1966
- No. 19 Severe Storm Warning Networks in Oklahoma. Gerald J. Carter and W. O. Garrison - July 1966

(continued) LIST OF SOUTHERN REGION TECHNICAL MEMORANDA

- No. 20 Climatological Aids to Short Range Forecasting. Jerome H. Codington - August 1966
- No. 21 A review of the Methods Developed for Forecasting Stratus in South Central Texas. Richard S. Schrag - August 1966
- No. 22 Agricultural Forecasting at Tallahassee, Florida. J. S. Smith - August 1966
- No. 23 The Use of Radar in Flash Flood Forecasting. Jack L. Teague - August 1966
- No. 24 Short Range Forecasting Procedures at Savannah. David P. Barnes - August 1966
- No. 25 On the Use of Digitized Radar Data for the Florida Peninsula. Neil L. Frank Paul L. Moore and George E. Fisher - Sept. 1966
- No. 26 Relative Humidity in Georgia. Horace S. Carter - Sept. 1966
- No. 27 Study on Duration of measurable Precipitation at Birmingham. Hugh B. Riley - September 1966
- No. 28 The Weather Distribution with Upper Tropospheric Cold Lows in the Tropics. Neil L. Frank - September 1966
- No. 29 On the Correlation of Radar Echoes over Florida with Various Meteorological Parameters. Neil L. Frank and Daniel L. Smith - October 1966
- No. 30 A study of the Diurnal Summer Wind System at Galveston, Texas. David H. George - December 1966
- No. 31 A simple Inexpensive Degree-Hour Counter. D. R. Davis and Jerrell E. Hughes - March 1967
- No. 32 An Objective Technique for Forecasting the Possibility of an Afternoon Summer Shower at Savannah, Georgia. David P. Barnes, Jr. and Samuel C. Davis - March 1967
- No. 33 An Empirical Method for Forecasting Radiation Temperatures in the Lower Rio Grande Valley of Texas. Leroy B. Hagood - March 1967
- No. 34 Study on Duration of Measurable Precipitation at Lubbock, Texas. G. Alan Johnson and Thomas P. Clarke - April 1967
- No. 35 Remoting Radar Scope Weather and Associated Data via the Slo-Scan Method. Davis Benton - May 1967
- No. 36 Short Range Forecasting of Dryoff Time From Dew Block Dew Intensity. Dorus D. Alderman and Kenneth E. Bryan - October 1967